

Dynamics of few-body states in a medium

M. Beyer*, S. Mattiello*, S. Strauß*, T. Frederico^{†,*}, H. J. Weber**,
P. Schuck[‡] and S.A. Sofianos[§]

**Fachbereich Physik, Universität Rostock, D-18051 Rostock, Germany*

[†]*Dep. de Física, ITA, CTA, 12.228-900 São José dos Campos, São Paulo, Brazil*

***Dept. of Physics, University of Virginia, Charlottesville VA, U.S.A.*

[‡]*Institut de Physique Nucléaire, F-91406, Orsay Cedex, France*

[§]*Physics Department, University of South Africa, Pretoria 0003, South Africa*

Abstract. Strongly interacting matter such as nuclear or quark matter leads to few-body bound states and correlations of the constituents. As a consequence quantum chromodynamics has a rich phase structure with spontaneous symmetry breaking, superconductivity, condensates of different kinds. All this appears in many astrophysical scenarios. Among them is the formation of hadrons during the early stage of the Universe, the structure of a neutron star, the formation of nuclei during a supernova explosion. Some of these extreme conditions can be simulated in heavy ion colliders. To treat such a hot and dense system we use the Green function formalism of many-body theory. It turns out that a systematic Dyson expansion of the Green functions leads to modified few-body equations that are capable to describe phase transitions, condensates, cluster formation and more. These equations include self energy corrections and Pauli blocking. We apply this method to nonrelativistic and relativistic matter. The latter one is treated on the light front. Because of the medium and the inevitable truncation of space, the few-body dynamics and states depend on the thermodynamic parameters of the medium.

Many-body Green functions. To treat hot and dense quantum systems, such as nuclear or quark matter in question here, we use the techniques of many-body Green functions [1]. We organize the Green functions into a Dyson expansion that leads to a hierarchy of linked cluster equations [2]. In recent years we have systematically applied this approach for finite temperatures up to multi-nucleon clusters (see e.g. [3] and refs therein) and also generalized it to light front dynamics [4, 5]. In fact, the light front quantization is well suited to use the many-body Green function formalism as in the relativistic case, as it utilizes the Fock space expansion [6]. We define a chronological Green function

$$\mathcal{G}_{\alpha\beta}^{t-t'} = -i \left(\theta(t-t') \langle A_{\beta}^{\dagger}(t') A_{\alpha}(t) \rangle \mp \theta(t'-t) \langle A_{\beta}^{\dagger}(t') A_{\alpha}(t) \rangle \right). \quad (1)$$

The average $\langle \dots \rangle$ is taken over the exact ground state and the upper (lower) sign is for fermions (bosons). The operators $A(t) = e^{itH} A e^{-itH}$ could be build out of any number of field operators (fermions and/or bosons). In finite temperature formalism the above definition can be generalized. For a grand canonical ensemble on the light front, the generalized Heisenberg picture assumes the form $A(\tau) = e^{\tau H} A e^{-\tau H}$, where $H = u \cdot P - \mu N$. Here $u, u^2 = 1$ is the four-velocity of the medium. For a nonrelativistic medium at rest $u^0 = 1, \vec{u} = 0$, hence $H = P_0 - \mu N$ with the Hamiltonian P_0 , suffice. On the light front $u^{\mu} = (u^+, u^-, \vec{u}_{\perp}) = (1, 1, 0, 0)$ [4]. The ensemble average is taken over the




FIGURE 1. Fraction of protons, neutrons (both identical in symmetric nuclear matter, long-dashed), deuteron (dots), triton, ${}^3\text{He}/{}^3\text{H}$ (both dashed), and α particles (solid) as a function of collision time at $T = 10$ MeV.




FIGURE 1. Phase diagram of QCD, chiral restoration (solid), pion dissociation line (dotted), nucleon dissociation lines for different cut-offs, $\Lambda = 4m$ (long dashed), $6m$ (dashed-dotted), $8m$ (dahed), m medium dependent quark mass.

(equilibrium) grand canonical statistical operator $\rho_G = Z_G / \text{Tr} Z_G$, viz. $\langle \dots \rangle = \text{tr} \{ \rho_G \dots \}$ and

$$Z_G = \exp \{ -(u \cdot P - \mu N) / T \}.$$

Dyson equations can be established for real time and imaginary time Green functions [2], also in light-front form [4]. Neglecting memory or retardation terms the equation for the Green function is given by

$$\left(i \frac{\partial}{\partial t} - \mathcal{M}_0^t \right) \mathcal{G}^{t-t'} = \delta(t-t') \mathcal{N}^t, \quad (2)$$

where $\mathcal{M}_{\alpha\beta}^t = \langle [[A_\alpha, H](t), A_\beta^\dagger(t)]_\pm \rangle$, $\mathcal{N}_{\alpha\beta}^t = \langle [A_\alpha, A_\beta^\dagger]_\pm(t) \rangle$. Formally, this equation holds for the nonrelativistic case and in light front form, with t replaced by the light front time $x^+ = t + z$ [4, 5]. Now it is possible by choosing the appropriate Fock operators, e.g. $A_\alpha = a_1 a_2 a_3$ for a three-body correlation, to derive from (2) cluster mean-field equations for the few-body system embedded in a medium [3, 4, 5].

Nuclear matter. At lower densities and temperatures nuclear matter is composed out of nucleons and nuclei. To describe heavy ion collisions at energies of about 30-100 MeV per nucleon it is sufficient to use a nonrelativistic treatment. Typical experiments have been performed by the INDRA collaboration [7]. At central collisions, where the system is likely to thermalize, the composition shows a large multiplicities of α -particles. This cannot be explained by a simple ideal gas of components. The nuclei forming the gas are influenced by the medium. The binding energies change and they can dissolve (Mott transition). The equations follow from the Dyson expansion as explained in the previous paragraph [3]. In a simple fire ball expansion (tailored to reproduce the Xe on Sn experiment at 50 MeV/A [7]) we assume a gas of nuclei expanding during the heavy ion collision. This is shown in Fig. 1. At around $t = 100 - 120$ fm/c the number of α -particles is much larger than the number of other clusters. This time,

which corresponds to about 3-4 times the initial volume, can be considered as freeze-out time. This freeze-out time essentially determines the composition of nuclei hitting the detector. This approach opens possibilities to address further questions related, e.g. to the influence of three-nucleon correlations on superconductivity, to four-nucleon condensation and even to larger clusters than α -particles.

Quark matter - light front form. At higher densities or temperatures we need a relativistic treatment. To this end we have generalized the light-front quantization to statistical quantum mechanics [3, 4, 5], see also [8]. The light-front quantization is capable to treat the perturbative and the nonperturbative region on the same footing. It can be realized by using the Nambu Jona-Lasinio model as an example. This approach reproduces the well known chiral phase transition, given as solid line in Fig. 2. We have also calculated the pion (dotted) and the nucleon (several dashed lines) dissociation line for different cut-off parameters. One may consider these lines related to a confinement-deconfinement transition, since no bound states are possible above the respective lines. The approach provides the framework to tackle such intriguing questions how hadrons form during the early plasma phase of the universe, if there is a color superconducting phase, what is the nature of QCD phase transition. It is also possible (and necessary) to go beyond the zero range model and implement real light front QCD, which has been elaborated in the works by Brodsky, Pauli and Pinsky [6] and is still progressing. The results are promising and since we do have a light front form of QCD available [6] it is exciting and challenging to extend this theory to quantum statistics.

Acknowledgments. We would like to thank the organizers for a pleasant and inspiring atmosphere during this meeting. Work supported by Deutsche Forschungsgemeinschaft.

REFERENCES

1. Fetter A L and Walecka J D 1971, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York), and Dover reprints 2003.
2. J. Dukelsky, G. Röpke, and P. Schuck Nucl. Phys. **A 628** (1998) 17 and refs. therein.
3. M. Beyer, G. Röpke and A. Sedrakian, Phys. Lett. B **376** (1996) 7; M. Beyer, W. Schadow, C. Kuhrts and G. Röpke, Phys. Rev. C **60** (1999) 034004; M. Beyer, S. A. Sofianos, C. Kuhrts, G. Röpke and P. Schuck, Phys. Lett. B **488** (2000) 247; M. Beyer, S. Strauss, P. Schuck and S. A. Sofianos, Eur. J. Phys. in print, arXiv:nucl-th/0310055; S. A. Sofianos and M. Beyer, arXiv:nucl-th/0408073.
4. M. Beyer, S. Mattiello, T. Frederico and H. J. Weber, Phys. Lett. B **521**, 33 (2001).
5. M. Beyer, S. Mattiello, T. Frederico and H. J. Weber Few Body Syst. **31** (2002) 159; M. Beyer, in Bled Workshops in Physics, eds. B. Golli, M. Rosina, S. Sirca, (Ljubljana 2003) p.11, arXiv:hep-ph/0310324; M. Beyer, S. Mattiello, T. Frederico and H. J. Weber, arXiv:hep-ph/0310222.
6. S. J. Brodsky, H. C. Pauli and S. S. Pinsky, Phys. Rept. **301**, 299 (1998), and refs. therein.
7. INDRA Collaboration, R. Nebauer *et al.*, Nucl. Phys. A658, 67 (1999); INDRA Collaboration, D. Gorio *et al.*, Eur. Phys. J. A **7**, 245 (2000), and references therein; INDRA Collaboration, S. Hudan *et al.*, Phys. Rev. C **67**, 064613 (2003). INDRA Collaboration, B. Borderie *et al.*, Phys. Lett. B **388**, 224 (1996); INDRA Collaboration, B. Borderie *et al.*, Phys. Lett. B **353**, 27 (1995).
8. V. S. Alves, A. Das and S. Perez, Phys. Rev. D **66** (2002) 125008; B. Blankleider and A. N. Kvinikhidze, Phys. Rev. D **69** (2004) 128701; J. Raufeisen and S. J. Brodsky, arXiv:hep-th/0408108.